

Regularity and finite injective dimension in characteristic $p > 0$

Tiberiu Dumitrescu and Cristodor Ionescu*

Faculty of Mathematics, University of Bucharest,
14 Academiei St., Bucharest, RO-010014, Romania
email tiberiu@al.math.unibuc.ro

Institute of Mathematics Simion Stoilow of the Romanian Academy,
P.O. Box 1-764, Bucharest, RO-014700, Romania
email Cristodor.Ionescu@imar.ro

Abstract

Recently, the regular local rings of prime characteristic were characterized in terms of the finiteness of injective dimension of the Frobenius map. We obtain relative versions of this result.

Let A be a noetherian local ring of prime characteristic $p > 0$ and let F_A denote the Frobenius morphism of A . A celebrated theorem of Kunz [5] asserts that A is regular if and only if F_A is flat. Later, Rodicio [9] has shown that A is regular if and only if F_A has finite flat dimension. Recently Avramov, Iyengar and Miller [2] have shown that A is regular if and only if F_A has finite injective dimension.

André and Radu [1],[8] proved a relative version of Kunz's result. In order to state it, we consider the following setup. For a ring C of prime characteristic $p > 0$, denote by $C^{(p)}$ the C -algebra structure on C given by F_C . Now let $u : A \rightarrow B$ be a morphism of noetherian rings of prime characteristic $p > 0$. We get the commutative diagram

*This paper was written while the authors were visiting the *School of Mathematical Sciences*, Lahore. They thank this institution for support and hospitality.

$$\begin{array}{ccc}
A & \xrightarrow{u} & B \\
F_A \downarrow & & \downarrow F_A \otimes_A B \\
A^{(p)} & \xrightarrow{u \otimes_A A^{(p)}} & A^{(p)} \otimes_A B \xrightarrow{w_{B/A}} B^{(p)}
\end{array}$$

where $w_{B/A}$ is given by $w_{B/A}(a \otimes b) = ab^p$, for every $a \in A$, $b \in B$. Note that $w_{B/A}$ induces a bijection on the spectra. In particular, if $w_{B/A}$ is flat and $B^{(p)}$ is noetherian, then $A^{(p)} \otimes_A B$ is also noetherian. The following result was proved by André, Radu and the first author of the present paper. See [1] and [8] for the equivalence of (a) and (b), and [4, Th. 2.13] for (c).

Theorem 1 *Let $u : A \rightarrow B$ be a morphism of noetherian rings. The following assertions are equivalent:*

- (a) *u is regular;*
- (b) *$w_{B/A}$ is flat;*
- (c) *u is flat and $w_{B/A}$ has finite flat dimension.*

The aim of this note is to investigate relative versions of the result of Avramov, Iyengar and Miller described above, in the spirit of Theorem 1.

All rings considered in this note have prime characteristic p . Notations and terminology are as in [7]. Consider the above setup. If the ring $A^{(p)} \otimes_A B$ is noetherian, from the previous results we obtain:

Theorem 2 *Let $u : A \rightarrow B$ be a flat local morphism of noetherian local rings. Suppose that $A^{(p)} \otimes_A B$ is noetherian. Then $w_{B/A}$ has finite injective dimension if and only if u is regular and A is Gorenstein.*

Proof. If $w_{B/A}$ has finite injective dimension, [2, Th. 13.2] implies that $A^{(p)} \otimes_A B$ is Gorenstein and $w_{B/A}$ has finite flat dimension. Applying Theorem 1, we obtain that u is regular. Since u is faithfully flat, $A^{(p)} \otimes_A B$ is also faithfully flat over $A^{(p)}$. Hence A is Gorenstein.

Conversely, since u is regular and A is Gorenstein, B is also Gorenstein. By Theorem 1 it follows that $w_{B/A}$ is faithfully flat, hence $A^{(p)} \otimes_A B$ is Gorenstein. [2, Th. 13.2] shows that $w_{B/A}$ has finite injective dimension. •

Corollary 3 *Let A be a noetherian local ring with geometrically regular formal fibers. Then A is Gorenstein if and only if $\widehat{A}$ has finite injective dimension over $A[\widehat{A}^p]$.*

Proof. From the assumption and Theorem 1 it follows that $w_{\widehat{A}/A}$ is flat, hence injective. Then the image of $w_{\widehat{A}/A}$ is $A[\widehat{A}^p]$. Now apply Theorem 2. •

Remark 4 Consider the diagram before Theorem 1. Let A' be an A -algebra and set $B' = B \otimes_A A'$. Then it is easy to check that $w_{B'/A'} = w_{B/A} \otimes_{A^{(p)}} A'^{(p)}$.

According to [4, Def. 2.8] a local ring is called *quasi-basic* if the A -algebra $A^{(p)}$ is a directed union of a family $(A_i)_{i \in I}$ of finite subalgebras, such that $A^{(p)}$ is flat over A_i , for any $i \in I$. By [4], any ring with Artin's approximation property (e.g. a complete local ring) is quasi-basic. Also, by [4, Prop. 2.9], any algebra essentially of finite type over a quasi-basic ring is quasi-basic. In particular, algebras which are essentially of finite type over a field are quasi-basic. Denote by $\text{fd}_A(M)$ (resp. $\text{id}_A(M)$), the flat (resp. injective) dimension of an A -module M .

Lemma 5 *Let $f : R \rightarrow S$ be a local morphism of local noetherian rings such that $\text{id}_R(S) < \infty$. Then $\text{fd}_R(S) \leq \text{id}_R(S) + \text{edim}(S)$.*

Proof. Consider a minimal Cohen factorization of the canonical morphism $R \rightarrow \widehat{S}$ (see [2])

$$\begin{array}{ccc} R & \longrightarrow & S \\ \downarrow & & \downarrow \\ R' & \longrightarrow & \widehat{S}. \end{array}$$

From [2, Cor. 2.5] we get

$$\text{id}_{R'}(\widehat{S}) \leq \text{id}_R(S) + \text{edim}(S) < \infty.$$

By [2, Th. 13.2], R' is a Gorenstein ring. From [6, Th. 2.2] we obtain $\text{fd}_{R'}(\widehat{S}) < \infty$ so, by Auslander-Buchsbaum formula,

$$\text{fd}_{R'}(\widehat{S}) \leq \text{depth}(R') = \text{id}_{R'}(\widehat{S}).$$

So, from [2, Cor. 2.5] we obtain

$$\text{fd}_R(S) \leq \text{fd}_{R'}(\widehat{S}) \leq \text{id}_{R'}(\widehat{S}) \leq \text{id}_R(S) + \text{edim}(S). \bullet$$

Theorem 6 *Let $u : A \rightarrow B$ be a local morphism of noetherian local rings. Assume that A is quasi-basic and $w_{B/A}$ has finite injective dimension. Then $w_{B/A}$ has finite flat dimension. Moreover if u is flat, then u is regular.*

Proof. Set $w := w_{B/A}$ and $C := A^{(p)} \otimes_A B$. Since A is a quasi-basic ring, the A -algebra $A^{(p)}$ is a directed union of a family $(A_i)_{i \in I}$ of finite subalgebras such that $A^{(p)}$ is flat over A_i , for any $i \in I$. Then $B_i := A_i \otimes_A B$ is a noetherian local ring. Consider the following commutative diagram with canonical morphisms:

$$\begin{array}{ccccc} A & \xrightarrow{u} & B & & \\ \tau_i \downarrow & & \downarrow v_i & & \\ A_i & \longrightarrow & B_i & & \\ \sigma_i \downarrow & & \downarrow s_i & & \\ A^{(p)} & \longrightarrow & C & \xrightarrow{w} & B^{(p)}. \end{array}$$

Since σ_i is flat, it follows that $s_i = \sigma_i \otimes_A B$ is faithfully flat. Hence

$$\mathrm{Ext}_C^n(E \otimes_{B_i} C, B^{(p)}) \cong \mathrm{Ext}_{B_i}^n(E, B^{(p)})$$

for any B_i -module E and for any $n \in \mathbb{N}$, according to [10, Th. 11.65]. It follows that $\mathrm{id}(ws_i) \leq \mathrm{id}(w)$. From Lemma 5 we have

$$\mathrm{fd}(ws_i) \leq \mathrm{id}(ws_i) + \mathrm{edim}(B) := k.$$

Let $j > k$. Then $\mathrm{Tor}_j^{B_i}(B^{(p)}, E) = 0$, for any B_i -module E . A direct limit argument (see [3, Ch. VI, Exercise 17]) shows that $\mathrm{Tor}_j^C(B^{(p)}, F) = 0$, for any C -module F . Thus $\mathrm{fd}(w) \leq k$. In addition, if u is flat, then u is regular by Theorem 1. •

Corollary 7 *Let $u : A \rightarrow B$ be a local morphism of noetherian local rings, such that A is a quasi-basic regular ring. If $w_{B/A}$ has finite injective dimension, then u is a regular morphism.*

Proof. Apply Theorem 6 and [4, Th. 2.10]. •

Let $u : A \rightarrow B$ be a morphism of noetherian local rings. It is well-known that when A is a field, u is a regular morphism if and only if B is a formally smooth A -algebra. From Theorems 2 and 6 we get the following consequence:

Corollary 8 *Let k be a field and B a noetherian local k -algebra. Then B is a formally smooth k -algebra if and only if $w_{B/k}$ has finite injective dimension. •*

When $k = \mathbf{Z}_p$, it follows that $w_{B/k} = F_B$, hence we retrieve [2, Th. 13.3]: A noetherian local ring B is regular if and only if F_B has finite injective dimension.

References

- [1] M. André, *Homomorphismes réguliers en caractéristique p* , C. R. Acad. Sci. Paris, **316** (1993), 643-646.
- [2] L. Avramov, S. Iyengar, C. Miller, *Homology over local homomorphisms*, Amer. J. Math., **128** (2006), 23-90.
- [3] H. Cartan, S. Eilenberg, *Homological Algebra*, Princeton Univ. Press, Princeton, New Jersey, 1993.
- [4] T. Dumitrescu, *Regularity and finite flat dimension in characteristic $p > 0$* , Commun. Algebra, **24** (1996), 3387-3401.
- [5] E. Kunz, *Characterizations of regular local rings of characteristic p* , Amer. J. Math., **41** (1969), 772-784 .
- [6] G. Levin, W. Vasconcelos, *Homological dimensions and Macaulay rings*, Pacific J. Math., **25** (1968), 315-323.
- [7] H. Matsumura, *Commutative Ring Theory*, Cambridge Univ. Press, Cambridge, 1986.
- [8] N. Radu, *Une classe d'anneaux noethériens*, Revue Roumaine Math. Pures Appl., **37** (1992), 79-82.
- [9] A. Rodicio, *On a result of Avramov*, Manuscripta Math., **62** (1988), 181-185.
- [10] J. Rotman, *An Introduction to Homological Algebra*, Academic Press, New York, 1979.